

Short communication

## *Erigeron acris* L. subsp. *angulosus* (Gaudin) Vacc. (Asteraceae), a new taxon in the flora of Poland

Artur Pliszko\*

Department of Taxonomy, Phytogeography and Paleobotany, Institute of Botany, Jagiellonian University, Kopernika 31, 31-501 Kraków, Poland

**Abstract** – The paper reports for the first time the occurrence of *Erigeron acris* subsp. *angulosus* in Poland. This rare European temperate plant was found in August 2014 in a former sand and gravel quarry, close to the Sobolewo reservoir in the town of Suwałki, north-eastern Poland. Species composition of the habitat is characterized by a phytosociological relevé based on the Braun-Blanquet method, diagnostic characters in comparison to the morphologically similar *E. acris* subsp. *droebachiensis* are presented using scanning electron microscopy imaging, and an identification key for *E. acris* s. l. in Poland is given.

**Keywords:** distribution, *Erigeron*, indumentum, Poland, scanning electron microscopy

### Introduction

*Erigeron acris* L. subsp. *angulosus* (Gaudin) Vacc. (Asteraceae), a European temperate representative of *E. acris* L. s. l. (Greuter 2003, 2006–2016), is confined mainly to the alpine zone of the Alps (Šída 1998, 2000, 2004). It has been reported from France, Italy, Switzerland, Liechtenstein, Austria, Slovenia, Germany, the Czech Republic, Slovakia, Hungary, Romania and Estonia (Greuter 2006–2016). This biennial or perennial herb usually occurs in scree and gravel river bank plant communities of the class *Thlaspietea rotundifolii* (Mucina 1997). One of the most reliable diagnostic characters that allows *E. acris* subsp. *angulosus* to be distinguished from most of the other subspecies of *E. acris* is the absence of villous indumentum on stems, leaves, and involucre bracts, with the exception of the short-stipitate glandular trichomes on involucre bracts (Šída 1998).

In Poland, a recent study based on herbarium materials of *E. acris* s. l. deposited in the Polish herbaria (Pliszko 2015) revealed the occurrence of three subspecies, namely *E. acris* subsp. *acris*, *E. acris* subsp. *droebachiensis* (O. F. Müll.) Arcang., and *E. acris* subsp. *serotimus* (Weihe) Greuter. The present paper provides the first record of *E. acris* subsp. *angulosus* in Poland and is aimed at showing the morphological characteristics of the taxon as well as its distribution and habitat.

### Materials and methods

*Erigeron acris* subsp. *angulosus* was identified based on morphological features given by Šída (1998, 2000, 2004) and its voucher specimens are deposited in the Herbarium of the Institute of Botany of the Jagiellonian University in Kraków (KRA). A phytosociological relevé followed the Braun-Blanquet method (Braun-Blanquet 1964). Names of syntaxa and diagnostic species followed Mucina (1997). Associated plants were identified after Wójciak (2003) and Rutkowski (2004). Names of taxa followed Mirek et al. (2002) and Ochrya et al. (2003). In morphometric and SEM studies *E. acris* subsp. *angulosus* was compared to the morphologically similar *E. acris* subsp. *droebachiensis*. Measurements were taken on herbarium specimens. Indumentum terminology followed Werker (2000). Specimens of *E. acris* subsp. *droebachiensis* used in the morphometric study were collected in Brzozowiec (south-eastern Poland) in July 2013 and 2014 (GPS coordinates: 49°25'54,36"N/22°11'57,36"E), whereas those used in the SEM study were collected in Toruń (north-central Poland) in July 2013 (GPS coordinates: 53°1'21,06"N/18°33'22,68"E). Samples of dry cauline leaves, peduncles, and capitula were mounted on aluminum stubs coated with double-sided conductive carbon tabs. Next, the samples were coated with gold in a sputter-coater and examined with a Hitachi S-4700 scanning electron microscope at

\* Corresponding author, e-mail: artur.pliszko@uj.edu.pl

an accelerating voltage of 10 kV, part of the standard procedure for viewing biological specimens in the SEM method (Bozzola and Russell 1999).

## Results and discussion

*Erigeron acris* subsp. *angulosus* was discovered in August 2014 on the south-eastern outskirts of the town of Suwałki, north-eastern Poland (GPS coordinates: 54°4'18,06"N/22°57'48,84"E; altitude: 157 m.a.s.l.). According to Zajac (1978), the locality is situated within the FB18 square unit of the ATPOL cartogram grid. A small population consisting of 25 individuals was found in a former sand and gravel quarry, close to the Sobolewo reservoir (Fig. 1). Vegetation in this anthropogenic habitat is represented by populations of grassland and ruderal plant species. The species composition in the site occupied by *E. acris* subsp. *angulosus* is characterized by the following phytosociological relevé: size of relevé: 25 m<sup>2</sup>; cover of herb layer: 20%; cover of moss layer: 40%; Ch.Cl. *Artemisietea vulgaris*: *Poa compressa* 2, *Melilotus albus* +; Ch.Cl. *Koelerio-Corynephoretea*: *Helichrysum arenarium* 1, *Arenaria serpyllifolia* +, *Herniaria glabra* +, *Ceratodon purpureus* (d) 3; Ch.Cl. *Festuco-Brometea*: *Artemisia campestris* 1, *Centaurea stoebe* +; others: *Erigeron acris* subsp. *angulosus* 1, *Hieracium pilosella* 1, *Oenothera* sp. 1, *Echium vulgare* +, *Erigeron acris* subsp. *acris* +, *Pinus sylvestris* +. Taking into consideration the selected quantitative features (Tab. 1), *E. acris* subsp. *angulosus* clearly shows shorter stems and synflorescences, and a lower number of capitula than *E. acris* subsp. *droebachiensis*. The number of cauline leaves is similar in both subspecies; however, the lower cauline leaves of *E. acris* subsp. *angulosus*

are much narrower than those of *E. acris* subsp. *droebachiensis*. Regarding the SEM micrographs (On-line Suppl. Figs. 1–3), in contrast to *E. acris* subsp. *droebachiensis*, *E. acris* subsp. *angulosus* has no unbranched multicellular uniseriate non-glandular trichomes on involucral bracts, peduncles, and cauline leaves. On the other hand, the unbranched short-stipitate multicellular biseriate glandular trichomes are densely distributed on involucral bracts and sparsely on peduncles in both subspecies. Moreover, involucral bracts in *E. acris* subsp. *angulosus* are serrated relatively more deeply at the apex than those in *E. acris* subsp. *droebachiensis* (On-line Suppl. Fig. 2).

The occurrence of *E. acris* subsp. *angulosus* in north-eastern Poland suggests that there may be a continuity of its geographical distribution between Central Europe and the Baltic states, since the taxon is present in Estonia (Greuter 2006–2016). Furthermore, it is worth mentioning that most of the specimens collected in Suwałki have relatively more cauline leaves than those characterized by Šída (1998, 2000, 2004). This can be explained by the fact that they were found flowering in late summer. *Erigeron acris* subsp. *angulosus* typically flowers in June, and at higher altitudes in July (Šída 2000, 2004). For example, summer specimens of *E. acris* subsp. *acris* often produce more cauline leaves than typical late spring specimens, especially after mowing or grazing (Pliszko 2015). In this connection, it should also be mentioned that similar specimens with numerous glabrous cauline leaves and a low number of capitula were collected between Miedzierza and Sielpia Wielka (south-central Poland) by Walas in September 1929 (KRA 0124518). These specimens were mistakenly identified as *E. acris* subsp. *droebachiensis* (Pliszko 2015). However, it is hard to decide if they belong to *E. acris* subsp. *angulosus* because their atypical features seem to be a result of mowing or grazing. The locality between Miedzierza and Sielpia Wielka needs to be re-examined and the identity of the plant should be confirmed in the flowering stage in June or July.

*E. acris* subsp. *angulosus* is the rarest subspecies of *E. acris* in Poland. Although the habitat in which the plant was discovered shows an anthropogenic character, it should be viewed as a native taxon. Its presence was confirmed in July 2016 with one flowering specimen. The expansion of trees and shrubs during secondary succession and recreational use of motorcycles are the main factors that can pose a threat to the *E. acris* subsp. *angulosus* population in Suwałki.



Fig. 1. *Erigeron acris* subsp. *angulosus* in Suwałki, north-eastern Poland: A – upper part of the plant; B – habitat (photos by A. Pliszko).

Tab. 1. Morphological comparison of the selected quantitative characters of *Erigeron acris* subsp. *angulosus* and *E. acris* subsp. *droebachiensis*. N – number of examined individuals. Numbers indicate mean value (out of brackets), minimum and maximum values (in brackets).

Character	<i>Erigeron acris</i> subsp. <i>angulosus</i> (N=15)	<i>Erigeron acris</i> subsp. <i>droebachiensis</i> (N=15)
height of stem (cm)	31.9 (22.0–45.0)	46.6 (28.5–62)
number of cauline leaves	17.4 (12–24)	19.2 (10–25)
size of lower cauline leaf (cm)	3.8 × 0.4 (2.5–6.5 × 0.3–0.6)	7.6 × 1.0 (4.5–12.0 × 0.7–1.4)
number of capitula	6.2 (3–9)	21.3 (6–46)
length of synflorescence (cm)	6.6 (4.5–10.0)	14.1 (7–24)

Identification key for *Erigeron acris* s. l. in Poland

1. Stems and leaves densely villous, involucre bracts abaxially densely villous and sparsely short-stipitate glandular ..... 2  
     Stems and leaves glabrous or sparsely villous at the base, involucre bracts abaxially densely short-stipitate glandular, glabrous or sparsely villous ..... 3
2. Cauline leaves 4–16, flat, usually erect; internodes 1.0–8.0 cm; capitula in paniculiform synflorescence ..... *Erigeron acris* subsp. *acris*
3. Basal leaves with acute apex; cauline leaves narrowly lanceolate to linear; peduncles sparsely short-stipitate glandular or glabrous; capitula in racemiform synflorescence ..... *Erigeron acris* subsp. *angulosus*  
     Basal leaves with obtuse apex; cauline leaves lanceolate; peduncles densely to sparsely villous and short-stipitate glandular; capitula in paniculiform synflorescence ..... *Erigeron acris* subsp. *droebachiensis*

## Acknowledgements

I would like to thank M.Sc. Eng. Anna Łatkiewicz (Department of Mineralogy, Petrology and Geochemistry of the Institute of Geological Sciences of the Jagiellonian University in Kraków) for her help in taking SEM images. I also thank

the anonymous reviewers for their suggestions and comments. This work was financially supported by the Jagiellonian University in Kraków (grants for research projects DS/MND/WBiNoZ/IB/4/2013 and DS/MND/WBiNoZ/IB/2/2014).

## References

- Bozzola, J. J., Russell, L.D., 1999: Electron Microscopy. Principles and Techniques for Biologists. 2nd ed. Jones and Bartlett Publishers, Boston, Toronto, London, Singapoure.
- Braun-Blanquet, J., 1964: Pflanzensoziologie, Grundzüge der Vegetationskunde. 3rd ed. Springer-Verlag, Wien-New York.
- Greuter, W., 2003: The Euro+Med treatment of Astereae (Compositae) – generic concepts and required new names. Willdenowia 33, 45–47.
- Greuter, W., 2006–2016: Compositae (pro parte majore). In: Greuter, W., Raab-Straube, E. von (eds.), Compositae. Euro+Med Plantbase – the information resource for Euro-Mediterranean plant diversity. Retrieved January 27, 2016 from <http://ww2.bgbm.org>
- Mirek, Z., Piękoś-Mirkowa, H., Zajac, A., Zajac, M., 2002: Flowering plants and pteridophytes of Poland, a checklist. W. Szafer Institute of Botany, Polish Academy of Sciences, Kraków.
- Mucina, L., 1997: Conspectus of classes of European vegetation. Folia Geobotanica et Phytotaxonomica 32, 117–172.
- Ochyra, R., Żarnowiec, J., Bednarek-Ochyra, H., 2003: Census catalogue of Polish Mosses. W. Szafer Institute of Botany, Polish Academy of Sciences, Kraków.
- Pliszko, A., 2015: Taxonomic revision and distribution of *Erigeron acris* s. l. (Asteraceae) in Poland. Phytotaxa 208, 021–033.
- Rutkowski, L., 2004: A key for identification of vascular plants of lowland Poland (in Polish). Wydawnictwo Naukowe PWN, Warszawa.
- Šída, O., 1998: Taxonomic problems in *Erigeron* sect. *Trimorpha* (Compositae) in Eurasia. Preslia 70, 259–269.
- Šída, O., 2000: *Erigeron acris* agg. in the Czech Republic and Slovakia. Zprávy České botanické společnosti 35, 1–33 (in Czech).
- Šída, O., 2004: *Erigeron* L. In: Slavík, B., Štěpánková, J. (eds.), Flora of the Czech Republic vol. 7, 140–153. Academia, Praha (in Czech).
- Werker, E., 2000: Trichome diversity and development. In: Hallahan, D. L., Gray, J. C. (eds.), Advances in botanical research. Incorporating advances in plant pathology. Plant trichomes, vol. 31, 1–35. Academic Press, London.
- Wójciak, H., 2003: Lichens, bryophytes, pteridophytes. Flora of Poland. Multico Oficyna Wydawnicza, Warszawa (in Polish).
- Zajac, A., 1978: Atlas of distribution of vascular plants in Poland (ATPOL). Taxon 27, 481–484.